arXiv:0902.4675v1 [math-ph] 26 Feb 2009

A Convenient Alternative for Series Manipulation via the Translation Operator

D. J. Priour, Jr[1]
1. *Department of Physics, University of Missouri, Kansas City, MO 64110-2499*

(Dated: February 26, 2009)

We derive and discuss a technique for manipulating power series which is complementary to standard procedures. We begin with the translation operator, but we express the operator as an infinite product instead of expanding it as a series and we apply combinatorial arguments to generate the terms in the series in an efficient manner with a minimum of clutter and intermediate calculations. The method is effective for developing multivariate expansions, and may also be used to manipulate series, e.g. in operations where one must take the reciprocal of a power series or raise it to a power that may be fractional or irrational. In the case of two component perturbations, we obtain analytic expressions for the expansion coefficients. We use our technique to generate an electrostatic multipole expansion as a demonstration of its utility in producing coefficients of special functions such as the Legendre and Hermite polynomials.

PACS numbers: PACS numbers: 02.30.Lt, 02.30.Mv, 02.30.Gp

I. INTRODUCTION

Taylor series expansions are an important tool in the analysis of the effect of small perturbations where the development of a power series in the perturbing influence conveniently organizes successive corrections to the zeroth order result. If a single variable function is written as $f(x_0+\Delta x)$, where Δx is the perturbation of small displacement, then the Taylor has the form

$$f(x_0+\Delta x) = f(x_0)+\Delta x\frac{d}{dx}f(x_0)+\frac{1}{2!}\Delta x^2\frac{d^2}{dx^2}f(x_0)+\ldots \tag{1}$$

A salient example familiar to the modern physics student is the case of the Lorentz factor central to formulae from special relativity. The Lorentz factor appears in the expression for the total energy of a body in motion, $\gamma m_0 c^2$, where $\gamma = (1-\beta^2)^{-1/2}$, m_0 is the rest mass, and c is the speed of light; $\beta = v/c$ is the object's speed relative to the speed of light. With a single variable Taylor series expansion, we readily obtain the energy of the mass when it is at rest, the classical kinetic energy term, and the first relativistic correction. One has

$$E = \gamma m_0 c^2 = (1-\beta^2)^{-1/2} m_0 c^2 \tag{2}$$

For ordinary speeds (i.e. for very small β), the Lorentz factor can be expanded as

$$\gamma = (1-\beta^2)^{-1/2} = 1+\frac{1}{2}\beta^2+\frac{3}{4}\beta^4+\ldots \tag{3}$$

Thus, we find for the total energy of the moving mass

$$E = \gamma m_0 c^2 = m_0c^2+\frac{1}{2}m_0v^2+\frac{3}{4}m_0v^4/c^4 \tag{4}$$

$$= m_0c^2+\frac{1}{2}m_0v^2\left(1+\frac{3}{2}\beta^2\right), \tag{5}$$

with the kinetic energy being augmented by the factor $1+\frac{3}{2}\beta^2$.The latter correction becomes more important at relativistic speeds where the kinetic energy scales more rapidly than the Newtonian quadratic dependence.

For situations involving more than one perturbing influence such as the function $f(x_0+\Delta x, y_0+\Delta y, z_0+\Delta z)$ where there are several displacements, a multivariable Taylor series expansion is often appropriate. As an example, one could have an expression raised to a fractional power such as $(1+[\Delta x+\Delta y+\Delta z])^{\alpha}$ where the perturbations are combined in a single term; in fact we will see that broadly speaking, the analysis of expressions of the form $f(1+[\Delta x+\Delta y+\Delta z])$ will be facilitated with the techniques we report on here. To develop the power series in the displacements, one could begin with

$$f[1+(\Delta x+\Delta y+\Delta z)] = 1+f^{'}(1)[\Delta x+\Delta y+\Delta z] + \frac{1}{2!}f^{''}(1)[\Delta x+\Delta y+\Delta z]^2+\ldots \tag{6}$$

From Eq. 6, we see that we have first the task of generating the primary series and then the evaluation of the individual terms, which will be polynomials (i.e. binomials for the case of two variables, trinomials for the three variable case, etc.) raised to successively increasing integer powers.

The case of a perturbation which happens to consist of a power series expansion in a small parameter can also be a computationally intensive scenario if one wishes to calculate a perturbative expansion in the small quantity. In the discussion which follows, we will use x as a generic label for the latter. As an example, one may consider $\sec(x) = 1/\cos(x)$, where

$$\sec(x) = (1-x^2/2!+x^4/4!-x^6/6!+\ldots)^{-1} \tag{7}$$

$$= 1-(-x^2/2!+x^4/4!-x^6/6!+\ldots)+ \tag{8}$$

$$(-x^2/2!+x^4/4!-x^6/6!+\ldots)^2+\ldots \tag{9}$$

Again, even after the initial expansion, the individual terms will need to be evaluated and like powers of x combined.

We now discuss a complementary and less laborious approach for the development of a Taylor series where the perturbation has a composite form such as the multivariable case in Eq. 6 or the situation where the perturbing influence is itself a power series in a smaller parameter, x. Our technique is based on the translation operator which can induce a coordinate shift by generating a Taylor series in one or more variable, and the method eliminates intermediate calculations while minimizing clutter.

II. THE SINGLE VARIABLE CASE

We comment first on the single variable case; excellent discussions may be found in standard texts[1]. Consider a function $f(x_0+\Delta x)$ where as before there is a shift about x_0 of Δx. For infinitesimal translations δx, reasonably behaved functions can be accurately rendered in linearized form with $f(x_0+\delta x) = f(x_0) + \delta x f^{'}(x_0)$. In terms of the differential operator $\hat{d}/dx$ one has, equivalently,

$$f(x_0+\delta x) = (1+\delta x\hat{d}/dx) f(x)|_{x=x_0} \tag{10}$$

It is possible to build up several shifts with

$$f(x_0+n\delta x) = (1+\delta x\hat{d}/dx)^n f(x)|_{x=x_0} \tag{11}$$

where there are n copies of the differential translation operator $(1+\delta\hat{d}/dx)$. This logic suggests that we may more accurately take into a finite shift Δx by combining many smaller translations to set up the larger shift. Hence, it is reasonable to suggest that

$$f(x_0+\Delta x) = \lim_{N\to\infty}(1+\Delta x/N)^N \tag{12}$$

becomes an increasingly accurate approximation as N becomes very large. The identity $\lim\limits_{N\to\infty}(1+\Delta x/N)^N = e^{\Delta x}$ suggests that we associate a finite translation Δx with the action of the exponentiated differential operator and well known result

$$\hat{T}(\Delta x) = e^{\Delta x\hat{d}/dx} \equiv \lim_{N\to\infty}(1+\Delta x/N)^N \tag{13}$$

In conjunction with this result, we apply combinatorial logic to develop a series in $\Delta x\hat{d}/dx$. Although the series will turn out to be nothing other than the standard single variable Taylor expansion given in Eq 1, a very similar approach will be applied to obtain the terms of the multivariable expansion where there are more intricacies to navigate. Returning to the infinite product form, $(1+\Delta x/N\hat{d}/dx)^N$, we examine how the power series will take shape as N becomes very large.

The zeroth order term in the power series will clearly be unity. To obtain the first order term, we must mix in a factor of $\frac{\Delta x}{N}\frac{\hat{d}}{dx}$, and there are a total of N ways of doing this. For the second order term, there are N ways to choose the first factor of $\frac{\Delta x}{N}\frac{\hat{d}}{dx}$, and $N-1$ ways to choose the second factor. However, with the order not being relevant, there will be redundant terms, and we must divide by a permutation factor of 2! to compensate for double counting. Proceeding in a similar way, we see that the prefactor for the third order term will be $N(N-1)(N-2)/3!$. Combining these results, we see that the entire series will have the form

$$(1+\frac{\Delta x}{N})^N f(x) = 1+\Delta x\frac{N}{N}\frac{df}{dx}+\Delta x^2\frac{N(N-1)}{2!N^2}\frac{d^2 f}{dx^2}+\ldots \tag{14}$$

Finally, in the large N limit, factors involving N cancel and we obtain the anticipated result

$$\left(1+\frac{\Delta x}{N}\frac{\hat{d}}{dx}\right)^N = 1+\Delta x\frac{df}{dx}+\frac{1}{2!}\Delta x^2\frac{d^2}{dx^2}+\ldots \tag{15}$$

III. THE MULTIVARIABLE CASE

For the multivariable case $f(y_1+\Delta_1, y_2+\Delta_2,\ldots)$, we will effect the translation with

$$\hat{T}(\Delta_1,\Delta_2,\ldots,\Delta_n) = \tag{16}$$

$$\exp\left(\Delta_1\frac{\hat{\partial}}{\partial y_1}+\Delta_2\frac{\hat{\partial}}{\partial y_2}+\ldots+\Delta_n\frac{\hat{\partial}}{\partial y_n}\right) \tag{17}$$

Noting that $\Delta_1\frac{\hat{\partial}}{\partial y_1}+\Delta_2\frac{\hat{\partial}}{\partial y_2}+\ldots+\Delta_n\frac{\hat{\partial}}{\partial y_n} = \vec{\Delta y}\cdot\vec{\hat{\nabla}}$, this becomes

$$\hat{T}(\Delta_1,\Delta_2,\ldots,\Delta_n) = \exp\left(\vec{\Delta y}\cdot\vec{\hat{\nabla}}\right) \tag{18}$$

En route to the multivariable Taylor series, one might develop a power series in $\vec{\Delta y}\cdot\vec{\hat{\nabla}}$, but the result will be a multivariable Taylor series requiring the subsequent evaluation of terms with polynomials raised to integer powers as discussed earlier. Instead, just as we did for the single variable case, we write the multivariable expansion as an infinite product in lieu of an infinite series. The result is

$$\hat{T}(\Delta_1,\Delta_2,\ldots,\Delta_n) = \tag{19}$$

$$\left(1+\frac{1}{N}\left[\Delta_1\frac{\hat{\partial}}{\partial y_1}+\Delta_2\frac{\hat{\partial}}{\partial y_2}+\ldots+\Delta_n\frac{\hat{\partial}}{\partial y_n}\right]\right)^N, \tag{20}$$

where it is understood that the large N limit will be taken.

Using combinatorial logic in conjunction with Eq. 20 will permit the immediate enumeration of specific terms. As an example, the coefficient for the linear term Δ_i will be $\Delta_i\frac{N}{N}\frac{\partial}{\partial y_i}f(y_1,y_2,\ldots y_n)$ with the N factors canceling. For quadratic terms, there are two cases. When $i\neq j$ there are $N(N-1)$ ways of collecting terms with the factors $\Delta_i\Delta_j$; we obtain $\Delta_i\Delta_j\frac{\partial^2}{\partial y_i\partial y_j}f(y_1,y_2,\ldots)$ in the large N limit. Note that since i and j are distinct, there is no need to divide by a redundancy factor. On the other hand, when $i=j$, we need to compensate for double counting by dividing by 2!, and the term is $\frac{1}{2!}\Delta_i^2\frac{\partial^2}{\partial y_i^2}f(y_1,y_2,\ldots y_n)$ as $N\to\infty$.

In general, one obtains a partial derivative $\partial/\partial y_j$ for each copy of Δ_j, and we must divide by the permutation factor $n_j!$ to compensate for multiple counting. As an example, the coefficient of the term $\Delta_1\Delta_2^2\Delta_3\Delta_4^4$ would be $\frac{1}{2!}\frac{1}{4!}\frac{\partial^8}{\partial y_1 \partial y_2^2 \partial y_3 \partial y_4^4} f(y_1, y_2, y_3, y_4, \ldots, y_n)$.

A salient and frequently occurring case where we will gain an advantage are situations in which the perturbing elements are combined together in a single term, i.e. as in $f(1 + [\Delta_1 + \Delta_2 + \ldots + \Delta_n])$. An important example is the case of an expression raised to a power α which needn't be an integer (though there are nontrivial cases that are often of interest such as the reciprocal, where $\alpha = -1$). Our expression is hence

$$f(y_1 + \Delta_1, y_2 + \Delta_2, \ldots, y_n + \Delta_n) \tag{21}$$

$$= (1 + [\Delta_1 + \Delta_2 + \ldots + \Delta_n])^\alpha \tag{22}$$

What will work in our favor and facilitate the calculation both in this special case and the broader class of functions $f(1+[\Delta_1+\Delta_2+\ldots+\Delta_n])$ is the fact that the partial derivatives depend only on the total order of the derivative, not the variable with which one differentiates. As an example, the derivatives $\frac{\partial^3}{\partial y_1 \partial y_2 \partial y_3} f(y_1, y_2, y_3, \ldots, y_n) = \frac{\partial^3}{\partial y_4^3} f(y_1, y_2, y_3, \ldots, y_n) = \alpha(\alpha-1)(\alpha-2)$ are identical. Let us examine the consequences of this simplifying property.

For linear terms, the partial derivative $\frac{\partial f}{\partial y_i}$ yields a factor α (independent of the value of i), and this value is the expansion coefficient. For the quadratic terms, the partial differentiation will yield for each $\{i, j\}$ the same factor $\alpha(\alpha - 1)$. In addition, we will need to divide by the permutation factor 2! if the two indices i and j are identical.

In general, as we continue to the third and higher order terms, the only things to keep in mind are the overall order of the term (the value of the derivative of f is fixed in this way), and the instances of repeated indices where it will be necessary to divide by a multiplicity factor for each case where there is a repeated index. For brevity, we represent the expansion series as $(1+[\Delta_1+\Delta_2+\ldots+\Delta_n])^\alpha = 1 + [A_1\Delta_1 + A_2\Delta_2 + \ldots + A_n\Delta_n] + [A_{11}\Delta_1^2 + A_{12}\Delta_1\Delta_2] + \ldots$ A few examples are

$$A_{12} = \alpha(\alpha-1);\; A_{11} = \frac{1}{2!}\alpha(\alpha-1) \tag{23}$$

$$A_{1222} = \frac{1}{3!}\alpha(\alpha-1)(\alpha-2)(\alpha-3) \tag{24}$$

$$A_{1234} = \alpha(\alpha-1)(\alpha-2)(\alpha-3) \tag{25}$$

Our results for the multivariable case also have relevance for perturbations which are themselves power series in a single variable. One may have, for example, an expression of the form

$$f(x) = (1 + [c_1 x + c_2 x^2 + c_3 x^3 + \ldots])^\alpha, \tag{26}$$

where the series may terminate or instead be an infinite power series. To make use of the machinery we have discussed in the case of more than one variable, we substitute a Δ_i for each term in the series (e.g. we set $\Delta_1 \equiv c_1 x$, $\Delta_2 = c_2 x^2$, etc). For convenience in sorting the terms, we choose the index i to correspond to the degree of the term it represents. As an example, in developing the power series for sec(x), we begin with

$$\sec(x) = 1/[1 + (-x^2/2! + x^4/4! - x^6/6! + \ldots)], \tag{27}$$

and for practical purposes that will become apparent we follow the pattern mentioned above and use the substitution $\sec(x) = 1/(1 + \Delta_2 + \Delta_4 + \Delta_6 + \ldots)$, where $\Delta_2 = -x^2/2!$, $\Delta_4 = x^4/4!$, and so on.

With the formalism we have discussed, we may now systematically and conveniently write down the terms in the expansion for $[1 + (\Delta_1 + \Delta_2 + \ldots)]^\alpha$, though we mention one caveat; it is important to list all possible combinations of the Δ_i factors in developing the expansion. For terms up to fourth order, this is easy to do by inspection. For the higher order cases, one can systematically enumerate all of the Δ_i combinations by listing all possible partitions of the integer corresponding to the order, and a recursive procedure for the partitioning is described in the Appendix. Since the value of the partial derivatives depends only on the total number of differentiations, we will use the notation d_i where, e.g., $d_1 = \partial f/\partial y_i$ and $d_2 = \partial^2 f/\partial y_i^2 = \partial^2 f/\partial y_i \partial y_j$ Following the rules derived earlier, the coefficients of powers of x are

$$d_1\Delta_1 = \alpha c_1 x$$

for the first order contribution,

$$d_1\Delta_2 + \frac{1}{2!}d_2\Delta_1^2 = \left[\alpha c_2 + \frac{1}{2!}\alpha(\alpha-1)c_1^2\right] x^2 \tag{28}$$

for the second order piece,

$$d_1\Delta_3 + d_2\Delta_2\Delta_1 + \frac{1}{3!}d_3\Delta_1^3 = \tag{29}$$

$$\left[\alpha c_3 + \alpha(\alpha-1)c_2c_1 + \frac{1}{3!}\alpha(\alpha-1)(\alpha-2)c_1^3\right] x^3$$

for the third order terms,

$$d_1\Delta_4 + d_2\Delta_3\Delta_1 + \frac{1}{2!}d_2\Delta_2^2 + \tag{30}$$

$$\frac{1}{2!}d_3\Delta_2\Delta_1^2 + \frac{1}{4!}d_4\Delta_1^4 =$$

$$\Big[\alpha c_4 + \alpha(\alpha-1)c_3c_1 + \frac{1}{2!}\alpha(\alpha-1)c_2^2 + \tag{31}$$

$$\frac{1}{2!}\alpha(\alpha-1)(\alpha-2)c_2c_1^2 + \frac{1}{4!}\alpha(\alpha-1)(\alpha-2)(\alpha-3)c_1^4\Big] x^4$$

for the fourth order contribution.

IV. THE TWO COMPONENT CASE AND A MULTIPOLE EXPANSION

A. Two component perturbations in general

We find a particularly simple result for the case of a two component perturbation, where in general we have

$f(1+[\Delta_1+\Delta_2])$ Since we will eventually replace $\Delta_1+\Delta_2$ with $c_1x+c_2x^2$ (i.e. as occurs often in generating functions for special functions such as the Legendre and Hermite polynomials), we will again separate the terms of the expansion into groups where the sum of the subscripts is the same; in this manner we will automatically have an appropriately organized power series in x when we make the substitution $y_1=c_1x$, $y_2=c_2x^2$. As is discussed in the Appendix, the fact that in this context one partitions integers into 1's and 2's with no higher integers appearing greatly limits the number of terms which will be generated for each order, and the number of terms associated with each order rises only linearly with the order n.

With these ideas in mind, we see that the first order contribution is $d_1\Delta_1$. The second order piece, with two terms, is $(d_2\Delta_2+\frac{1}{2!}d_2\Delta_1^2)$. The third order component also has two terms since there is no Δ_3 term in the perturbation, and we obtain $d_2\Delta_1\Delta_2+\frac{1}{3!}d_3\Delta_1^3$; likewise, there being no Δ_4 term, the fourth order piece has the form $d_2\Delta_2^2+\frac{1}{2!}d_3\Delta_2\Delta_1^2+\frac{1}{4!}d_4\Delta_1^4$, and one continues in this manner for the higher order terms. If we specialize to the case $y_1=c_1x$ and $y_2=c_2x^2$, we find for the power series in x

$$\begin{aligned} f(1+c_1x+c_2x^2) = f(1) + \Big\{ &\frac{1}{0!0!}d_1c_2^0c_1^1x + \\ &+\left(\frac{1}{1!1!}d_2c_2^1c_1^1+\frac{1}{0!3!}d_3c_2^0c_1^3\right)x^3\Big\} \\ &+\Big\{\left(\frac{1}{1!0!}d_1c_2^1c_1^0+\frac{1}{0!2!}d_2c_2^0c_1^2\right)x^2 \qquad (32) \\ +\left(\frac{1}{2!0!}d_2c_2^2c_1^0+\frac{1}{1!2!}d_3c_2^1c_1^2+\frac{1}{0!4!}d_4c_2^0c_1^4\right)&x^4+\ldots\Big\} \qquad (33) \end{aligned}$$

where 0!, 1!, c_2^0, and c_1^0 have been inserted strategically and odd and even terms separated to emphasize general trends. To obtain expressions for general even and odd powers, we combine the terms in parentheses into summations and thereby make the dependence on the order n more transparent. We deduce for the coefficients of x the expressions

$$\mathcal{A}_{2n}^{\text{even}} = \sum_{i=0}^{n}\frac{1}{(n-i)!2i!}d_{n+i}c_2^{n-i}c_1^{2i} \qquad (34)$$

$$\mathcal{A}_{2n-1}^{\text{odd}} = \sum_{i=1}^{n}\frac{1}{(n-i)!(2i-1)!}d_{n+i-1}c_2^{n-i}c_1^{2i-1} \qquad (35)$$

where we write for the full expansion $f(1+[c_1x+c_2x^2]) = f(1)+\mathcal{A}_1x+\mathcal{A}_2x^2+\mathcal{A}_3x^3+\ldots$

B. Multipole expansion for an electrostatic potential

A salient example from electrostatics where we may apply this logic is the multipole expansion of the electric potential of a charge displaced from the origin by Δz as shown in Figure 1. If we operate in the polar system, the combined potential due to the negative and positive charges is

$$\begin{aligned} V(r,\theta) = -Q/r[&(1-2\cos\theta\Delta/r+\Delta^2/r^2)^{-1/2} \\ &-(1+2\cos\theta\Delta/r+\Delta^2/r^2)^{-1/2}] \qquad (36) \end{aligned}$$

For distances large relative to the size of the charge arrangement (i.e. where $r\gg\Delta$), we may expand in the small parameter Δ/r.

The perturbation is a two term series in Δ/r; following the procedure described in this report, we set $(1+\Delta_1+\Delta_2)^{-1/2}\equiv(1-2\Delta/r\cos\theta+\Delta^2/r^2)^{-1/2}$ for the first term in the electrostatic potential. The analysis of the second term, except for a sign difference, proceeds in an identical way. Setting $c_1=-2\cos\theta$, $c_2=1$, and "x"$=\Delta/r$, we see that the first order contribution will be

$$d_1c_1x = -(1/2)(-2)\cos\theta(\Delta/r) = \cos\theta(\Delta/r), \qquad (37)$$

and this is $P_1(\cos\theta)(\Delta/r)$ where the $P_l(x)$ are Legendre polynomials of order l. The second order piece is

$$\begin{aligned} &\left(d_1c_2+\frac{1}{2!}d_2c_1^2\right)x^2 = \\ &[-1/2+(1/2)(-1/2)(-3/2)4\cos^2\theta](\Delta/r)^2 \\ &=(-1/2+3/2\cos^2\theta)(\Delta/r)^2 = P_2(\cos\theta)(\Delta/r)^2 \qquad (38) \end{aligned}$$

The third order contribution is

$$\begin{aligned} &\left(d_2c_1c_2+\frac{1}{3!}d_3c_1^3\right) = \\ &[(-1/2)(-3/2)(-2\cos\theta)+ \\ &(1/6)(-1/2)(-3/2)(-5/2)(-8\cos^3\theta)](\Delta/r)^3 = \\ &(-3/2\cos\theta+5/2\cos^3\theta)(\Delta/r)^3 = P_3(\cos\theta)(\Delta/r)^3 \qquad (39) \end{aligned}$$

The fourth and higher order terms are obtained in a similar manner. We have for the combined potential

$$V(r,\theta) = \frac{2\Delta Q}{r^2}\cos\theta+\frac{\Delta^3Q}{r^4}(-3\cos\theta+5\cos^3\theta)+\ldots \qquad (40)$$

We recognize the first nonzero term as the dipolar contribution where the factor $2\Delta Q$ is precisely the dipole moment appropriate to the geometry and the equal and opposite charges $+Q$ and $-Q$.

C. Additional Examples: Hermite Polynomials and the Bernoulli numbers

Generating functions often appear in the form $f[1+(c_1x+c_2x^2)]$, and we may therefore apply the formalism used in the context of the electric multipole expansion to generate the first several polynomials in such cases. For the Hermite polynomials, the generating function

$e^{tx-t^2/2}=g(x,t)^2$ is particularly amenable to our expansion technique since the derivatives $d_1=d_2=\ldots=d_n=1$ are each equal to unity. With $e^{tx-t^2/2}=\sum_{i=0}^{\infty}a_nt^n$ with $H_n(x)=n!a_n$, we find that $H_0(x)=1$ and $H_1(x)=(1!)(c_1)=x$, while $H_2(x)$ and $H_3(x)$ are given by

$$H_2(x)=2!\left(c_2+\frac{1}{2}c_1^2\right)=2\left(-\frac{1}{2}+\frac{x^2}{2}\right)=x^2-1, \tag{41}$$

and

$$H_3(x)=3!\left(c_2c_1+\frac{1}{3!}c_1^3\right)=6\left(-\frac{x}{2}+\frac{x^3}{6}\right)=x^3-3x. \tag{42}$$

Our technique can be used to conveniently obtain the Bernoulli numbers B_n from[3]

$$\frac{x}{e^x-1}=1+B_1x+\frac{B_2}{2!}x^2+\ldots \tag{43}$$

It is useful to write

$$\frac{x}{e^x-1}=\frac{1}{(1+x/2!+x^2/3!+\ldots)} \tag{44}$$

We see that $B_1=d_1c_1=-\frac{1}{2}$. For B_2, we have

$$B_2=2!\left(d_1c_2+\frac{1}{2!}d_2c_1^2\right)=2(-1/6+1/4)=\frac{1}{6} \tag{45}$$

We find for B_3

$$\begin{aligned}B_3&=3!\left(d_1c_3+d_2c_1c_2+\frac{1}{3!}d_3c_1^3\right)\\&=6(-1/24+1/6+1/8)=0\end{aligned} \tag{46}$$

D. Conclusions

In conclusion, we have discussed a technique for facilitating the manipulation of power series which relies on the structure of the multivariable translation operator when written as an infinite product. The technique eliminates intermediate calculations while minimizing clutter. The approach has been demonstrated for series raised to powers, but the technique may be applied in more general situations where one seeks to expand a function of the form $f[1+(a_1x+a_2x^2+\ldots)]$ where the perturbing influence is itself a power series in a small quantity x. In the case of a two component perturbation (e.g. as in generating functions for special functions such as the Legendre and Hermite polynomials), the process of generating the power series is relatively simple, and our technique facilitates the derivation of a analytical closed form expression for the coefficients of each order of x.

V. APPENDIX: THE PARTITIONING OF INTEGERS

The task of exhausting all possible combinations of Δy_i becomes increasingly subtle with increasing order. As an example, for the second order case one has only $\left\{\Delta_2,\Delta_1^2\right\}$, whereas for fifth order the set of possibilities is much larger, namely $\left\{\Delta_5,\Delta_4\Delta_1,\Delta_3\Delta_2,\Delta_3\Delta_1^2,\Delta_2^2\Delta_1,\Delta_2\Delta_1^3,\Delta_1^5\right\}$. Nevertheless, one can easily list all possibilities for small to moderate order by following a recursive partitioning procedure which we discuss here. We first examine the general case where the perturbing influence may have a large (or even infinite) number of terms. We then find a much simpler result for the case of a two component perturbation.

A. The general case

Let us denote the set of all possible partitions of the integer n with curly braces, (i.e. as $\{n\}$). It is easy to see that for $n=1$, there is only $\{1\}=[1]$.

Next, we discuss a systematic procedure for generating partitions by examining successively higher integers beginning with $n=2$ and continuing through $n=6$. One always begins with the unary partitioning [2], just the number itself enclosed in square brackets. The next step is to generate all unique binary partitions, and for $n=2$ there is one of these, [1, 1]. To represent the set of possible partitions, we use the group addition symbol $\oplus$ to indicate the combination of the individual partitions. E.G., for $n=2$, we have

$$\{2\}=[2]\oplus[1,1] \tag{47}$$

For $n=3$, we proceed in a similar way, again, forming all possible binary partitions and finding $\{3\}=[3]\oplus[\{2\},1]$. Now, substituting the partitioning $\{2\}$ determined for $n=2$, we obtain

$$\{3\}=[3]\oplus[[2]\oplus[1,1],1] \tag{48}$$

Exploiting linearity and distributing yields

$$\{3\}=[3]\oplus[2,1]\oplus[1,1,1] \tag{49}$$

for the partitioning corresponding to $n=3$.

For $n=4$, we have

$$\{4\}=[4]\oplus[\{3\},1]\oplus[\{2\},\{2\}] \tag{50}$$

Acting recursively and inserting the relations we obtained for $\{3\}$ and $\{2\}$ yields

$$\begin{aligned}\{4\}=[4]&\oplus[[3]\oplus[2,1]\oplus[1,1,1],1]\\&\oplus[[2]\oplus[1,1],[2]\oplus[1,1]]\end{aligned} \tag{51}$$

Distributing and eliminating redundant terms yields

$$\{4\}=[4]\oplus[3,1]\oplus[2,2]\oplus[2,1,1]\oplus[1,1,1,1] \tag{52}$$

We may proceed in the same manner for $n = 5$ and higher, but the structure of the partitionings $\{2\}$, $\{3\}$, and $\{4\}$ suggests a more efficient and straightforward approach akin to counting. To emphasize the counting character of the partitioning process, we have taken care to rank the specific partitions in decreasing "dictionary" order, where the first entry in the partitioning determines priority and ties are broken by subsequent entries where needed. We illustrate the systematic counting process for the case $n = 6$, and we obtain $\{6\}$ in this manner with no discarded terms.

The highest possible number which may appear in $\{6\}$ is just a single "6", so the first entry in the partitioning is $[6]$. The next highest entry would be a "5", and the only possible partition is $[5, 1]$. The highest ranking entry in the next tier is $[4, 2]$, and we may decrement by breaking up the lowest reducible entry; we hence split the "2" into two "1"'s yielding $[4, 1, 1]$. For the next level, the highest ranking entry is $[3, 3]$. The only way to generate a lower entry is to reduce the second "3", yielding $[3, 2, 1]$, followed by $[3, 1, 1, 1]$. The next level and the final tier will consist of the entries $[2, 2, 2] \oplus [2, 2, 1, 1] \oplus [2, 1, 1, 1, 1] \oplus [1, 1, 1, 1, 1, 1]$. To further illustrate the decrementation method, we generate the partitioning for $n = 8$ by counting down from $[8]$.

$$\begin{gathered}
\{8\} = [8] \oplus [7, 1] \\
\oplus [6, 2] \oplus [6, 1, 1] \\
\oplus [5, 3] \oplus [5, 2, 1] \oplus [5, 1, 1, 1] \\
\oplus [4, 4] \oplus [4, 3, 1] \oplus [4, 2, 2] \oplus [4, 2, 1, 1] \oplus [4, 1, 1, 1, 1] \\
\oplus [3, 3, 2] \oplus [3, 2, 2, 1] \oplus [3, 2, 1, 1, 1] \oplus [3, 1, 1, 1, 1, 1] \\
\oplus [2, 2, 2, 2] \oplus [2, 2, 2, 1, 1] \oplus [2, 2, 1, 1, 1, 1] \oplus [2, 1, 1, 1, 1, 1, 1] \\
\oplus [1, 1, 1, 1, 1, 1, 1, 1]
\end{gathered} \tag{53}$$

B. The case of a two component perturbation

With a two component perturbation such as $\Delta_1 + \Delta_2 = c_1 x + c_2 x^2$, the process of partitioning the integers for each order in x simplifies considerably, and can be done essentially by inspection for any order. As an example, we have for the first several orders

$$\begin{gathered}
\{1\} = [1]; \quad \{2\} = [2] \oplus [1, 1] \\
\{3\} = [2, 1] \oplus [1, 1, 1]; \quad \{4\} = [2, 2] \oplus [2, 1, 1] \oplus [1, 1, 1, 1]
\end{gathered}$$

---

[1] J. J. Safari, *Modern Quantum Mechanics, Revised Edition*, Addison-Wesley, Reading, Massachussetts (1994).

[2] Erwin Kreyszig, *Advanced Engineering Mathematics*, John Wiley and Sons, Inc., New York (1967).

[3] George Brown Arfken and Hans Jurgen Weber *Mathematical Methods for Physicists*,Sixth edition, Elsevier (2005).



Figure 1: The charge arrangement with Q and $-Q$ on the z axis and separated by a distance 2Δ.